


# Intact Stratigraphy Traversing the Phyllosilicate to Sulfate Eras at the Syrtis-Isidis Contact, Mars

*John Mustard, Bethany Ehlmann, J. R. Skok,  
Dave Des Marais, Nicolas Mangold, and  
Francois Poulet*


NOTE ADDED BY JPL WEBMASTER: This content has not been approved or adopted by, NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology






## Stratigraphy of Nili Fossae/NE Syrtis


record multiple aqueous environments from the Middle Noachian to Early Hesperian



**Stratigraphy of Nili Fossae/NE Syrtis**  
 record multiple aqueous environments from the Middle Noachian to Early Hesperian










# Chronology of Geological Processes Defined in a Mineralogic/Morphologic Stratigraphy Across 1000s km<sup>2</sup>




Isidis Basin and Syrtis  
Major lavas are major  
time-stratigraphic  
markers

Significant gradation  
(mass wasting? aeolian?  
alluvial? all?) between  
Isidis basin formation  
and Syrtis lava  
emplacement

Defined wet periods,  
marked by mineralogy &  
morphology




# The Noachian basement

- altered (to Fe/Mg smectite)
- breccia blocks (altered/unaltered) are common throughout
- raised ridges upon erosion:  
conduits of fluid flow?
- mafic (low-Ca pyx) materials cap
- Q: clays created or distributed by impact




- Low-Ca Pyroxene
- Fe/Mg smectites
- Fe-Phyllosilicate

*Mustard et al., 2009, JGR*







Widespread olivine-bearing unit is a distinctive stratigraphic marker  
Shows the same texture, thickness, and composition across 1000s km<sup>2</sup>



- Volcanic flows [Tornabene et al. 2007] vs impact melt [Mustard et al., 2007]
- Drapes topography and is cut by fossae
- Associated with carbonate and serpentine in the region near and north of Jezero Crater








## Morphology and Orientation of Ridges

Widespread ridges in the Noachian crustal unit, 10s m wide, 100s m long


Mapping of over 2200 ridges shows a NE-SW orientation that parallels Nili Fossae



## Morphology and Orientation of Ridges

Widespread ridges in the Noachian crustal unit, 10s m wide, 100s m long

Mapping of over 2200 ridges shows a NE-SW orientation that parallels Nili Fossae






# Fluvial Channels in Nili Fossae

Active from  
Noachian into  
Hesperian

Both well-  
developed  
drainage  
networks (e.g.  
Jezero) and  
short,  
amphitheatre  
headed valleys






## Kaolinite-smectite alteration


*occurs where precursor rock is not olivine-rich (pyx, Fe/Mg smectite)*

## Carbonate-smectite alteration

*occurs where precursor rock is olivine-rich*





Ehlmann et al., Science 2008;  
Ehlmann et al., JGR, 2009





# Kaolinite, carbonate, and serpentine mineralization

- Chemistry controlled by precursor mineralogy



- Carbonate: Surface alteration related to near-surface hydrology? Hydrothermal serpentinizing system?
  - $\text{H}_2$  an energy source for organisms, potential for methane production
- Kaolinite: Leaching from an active hydrologic system

## ***NE SYRTIS: Ellipse and Go-To Science***

*Cross the Noachian-Hesperian boundary and the transition from phyllosilicate/carbonate (alkaline) to sulfate (acidic)*


*Investigate hydrothermal, fluvial, and volcano-ice interactions which present a number of diverse habitable environments on early Mars*

*Thick sedimentary sequence is accessible at the go-to site*


***In-ellipse science includes immediate investigation of Noachian alteration in breccias and of kaolinite and carbonate alteration with reactants and products in direct association***


***Extended region includes four distinct aqueous environments with clear local stratigraphic relationships that fit within a regional stratigraphic and geologic framework***

***Fe/Mg smectite basement, kaolinite alteration, olivine-rich unit with serpentine and carbonate alteration, Hesperian volcanic flows emplaced on sediments and interacting with volatile-rich deposits with hydrothermal alteration and sulfate mineral deposition***













## 1. NE Syrtis inflow







Beneath the capping lava is a 500 m stack of layered rock that appears sedimentary






3. Beneath the capping lava is a 500 m stack of layered rock: Sedimentary?









## Regional Stratigraphy provides the context for in-ellipse and go-to science



## In Ellipse Science and Notional Traverses












# NE Syrtis Acid-Alkaline Transition

*Cross the Noachian-Hesperian boundary and the transition from phyllosilicate/carbonate (alkaline) to sulfate (acidic)*

*Emplacement of lava into a volatile-rich environment and onto the olivine-bearing units and Fe/Mg phyllosilicate-bearing basement with serpentine formation*

*Sulfate deposits include jarosite and polyhydrated sulfate, largely with volcanic units*

*Evidence for circulation of fluids by heat of lava source*



**Altered Noachian ultramafic unit**

*In direct stratigraphic section and in situ are units that encompass many elements of the transition from the early phyllosilicate-forming era to the later sulphate-forming era*

*Long record of aqueous processes with alteration minerals*

HRL0000B8C2 (R: 2.38  $\mu$ m, G: 1.80  $\mu$ m, B: 1.15  $\mu$ m)



# Conclusions

- Target-rich in ellipse science; go-to science traverses Noachian to Hesperian
- Bedrock strata in-situ representing four distinct environments of aqueous alteration where reactants and products are together
  - early crustal: creation or distribution by impact
  - carbonate/serpentine: surface alteration or hydrothermal?
  - layered phyllosilicates (Al- over Fe/Mg): from leaching with surface hydrology?
  - (sedimentary?) acid sulfate formation
- A record of aqueous geochemistry preserved in-situ, in mineral-bearing strata, distinct in age, primary mineralogy, and geologic setting well-suited for the MSL instrument suite
- Key stratigraphies from Bibring's Phyllosian and Theiikian eras: do the changes recorded here represent Mars global environmental change?